arXiv:1111.4071v1 [math.NT] 17 Nov 2011

Determinantal and permanental representation of generalized bivariate Fibonacci p-polynomials

Kenan KAYGISIZ[a,*], Adem SAHIN[a]

[a] *Department of Mathematics, Faculty of Arts and Sciences, Gaziosmanpaşa University, 60250 Tokat, Turkey*

Abstract

In this paper, we give some determinantal and permanental representations of generalized bivariate Fibonacci p-polynomials by using various Hessenberg matrices. The results that we obtained are important since generalized bivariate Fibonacci p-polynomials are general form of, for example, bivariate Fibonacci and Pell p-polynomials, second kind Chebyshev polynomials, bivariate Jacobsthal polynomials etc.

Keywords: Generalized bivariate Fibonacci p-polynomials, determinant, permanent and Hessenberg matrix.

---

1. Introduction

Fibonacci numbers and their generalizations have been studying for a long time. Miles [13] defined generalized order-k Fibonacci numbers(GOkF) as,

$$f_{k,n} = \sum_{j=1}^{k} f_{k,n-j} \tag{1}$$

for $n > k \geq 2$, with boundary conditions: $f_{k,1} = f_{k,2} = f_{k,3} = \cdots = f_{k,k-2} = 0$ and $f_{k,k-1} = f_{k,k} = 1$.

Er [2] defined k sequences of generalized order-k Fibonacci numbers ($k$SO$k$F) as; for $n > 0$, $1 \leq i \leq k$

$$f_{k,n}^{i} = \sum_{j=1}^{k} c_j f_{k,n-j}^{i}$$

with boundary conditions for $1 - k \leq n \leq 0$,

---

[*] Corresponding author (+903562521616-3087)

Email addresses: `kenan.kaygisiz@gop.edu.tr` (Kenan KAYGISIZ), `adem.sahin@gop.edu.tr` (Adem SAHIN)

$$f_{k,n}^{\,i} = \begin{cases} 1 & \text{if } \; i = 1-n, \\ 0 & \text{otherwise,} \end{cases}$$

where c_j $(1 \le j \le k)$ are constant coefficients, $f_{k,n}^{\,i}$ is the n-th term of i-th sequence of order-k generalization.

The generalized Fibonacci p-numbers [5] are

$$F_p(n) = F_p(n-1) + F_p(n-p-1) \tag{2}$$

for $n > p+1$, with boundary conditions $F_p(1) = F_p(2) = \cdots = F_p(p) = F_p(p+1) = 1$.

The Fibonacci [9], Pell [3], second kind Chebysev [17] and Jacobsthal [16] polynomial are defined as

$$\begin{aligned} f_{n+1}(x) &= xf_n(x) + f_{n-1}(x),\ n \ge 2 \text{ with } f_0(x) = 0,\ f_1(x) = 1 \\ P_{n+1}(x) &= 2xP_n(x) + P_{n-1}(x),\ n \ge 2 \text{ with } P_0(x) = 0,\ P_1(x) = x \\ U_{n+1}(x) &= xU_n(x) - U_{n-1}(x),\ n \ge 2 \text{ with } U_0(x) = 1,\ U_1(x) = 2x \\ J_{n+1}(x) &= J_n(x) + xJ_{n-1}(x),\ n \ge 2 \text{ with } J_0(x) = 0,\ J_1(x) = 1 \end{aligned}$$

respectively.

The generalized bivariate Fibonacci p-polynomials [16] are

$$F_{p,n}(x,y) = F_{p,n-1}(x,y) + F_{p,n-p-1}(x,y)$$

for $n > p$, with boundary conditions $F_{p,0}(x,y) = 0$, $F_{p,n}(x,y) = x^{n-1}$, $n = 1, 2, ..., p$.

A few terms of $F_{p,n}(x,y)$ for $p = 3$ and $p = 4$ are

$$\begin{gathered} 0, 1, x, x^2, x^3, y + x^4, 2xy + x^5, \ldots \\ 0, 1, x, x^2, x^3, x^4, y + x^5, 2xy + x^6, \ldots \end{gathered}$$

respectively.

MacHenry [10] defined generalized Fibonacci polynomials ($F_{k,n}(t)$) where t_i $(1 \le i \le k)$ are constant coefficients of the core polynomial

$$P(x; t_1, t_2, \ldots, t_k) = x^k - t_1 x^{k-1} - \cdots - t_k,$$

which is denoted by the vector $t = (t_1, t_2, \ldots, t_k)$.

$F_{k,n}(t)$ is defined inductively by

$$\begin{aligned} F_{k,n}(t) &= 0,\ n < 1 \\ F_{k,1}(t) &= 1 \\ F_{k,n+1}(t) &= t_1 F_{k,n}(t) + \cdots + t_k F_{k,n-k+1}(t). \end{aligned}$$

MacHenry studied on these polynomials and obtained very useful properties of these polynomials in [11, 12].

Remark 1.1. Pairs of cognate polynomial sequence [16]

x	y	p	$F_{p,n}(x,y)$
x	y	1	bivariate Fibonacci polynomials $F_n(x,y)$
x	1	p	Fibonacci $p-$polynomials $F_{p,n}(x)$
x	1	1	Fibonacci polynomials $f_n(x)$
1	1	p	Fibonacci $p-$numbers $F_p(n)$
1	1	1	Fibonacci numbers F_n
$2x$	y	p	bivariate Pell p-polynomials $F_{p,n}(2x,y)$
$2x$	y	1	bivariate Pell polynomials $F_n(2x,y)$
$2x$	1	p	Pell p-polynomials $P_{p,n}(x)$
$2x$	1	1	Pell polynomials $P_n(x)$
2	1	1	Pell numbers P_n
$2x$	-1	1	second kind Chebysev polynomials $U_{n-1}(x)$
x	$2y$	p	bivariate Jacobsthal p-polynomials $F_{p,n}(x,2y)$
x	$2y$	1	Bivariate Jacobsthal Polynomials $F_n(x,2y)$
1	$2y$	1	Jacobsthal Polynomials $J_n(y)$
1	2	1	Jacobsthal Numbers J_n

This Remark shows that $F_{p,n}(x,y)$ are general form of all sequences and polynomials mentioned in Remark 1.1. Therefore, any result obtained from $F_{p,n}(x,y)$ are valid for all sequences and polynomials mentioned in Remark 1.1.

Many researchers studied on determinantal and permanental representations of k sequences of generalized order-k Fibonacci and Lucas numbers. For example, Minc [14] defined an $n \times n$ (0,1)-matrix $F(n,k)$, and showed that the permanents of $F(n,k)$ is equal to the generalized order-k Fibonacci numbers (1).

The authors ([7], [8]) defined two $(0,1)$-matrices and showed that the permanents of these matrices are the generalized Fibonacci (1) and Lucas numbers. Öcal [15] gave some determinantal and permanental representations of k-generalized Fibonacci and Lucas numbers, and obtained Binet's formula for these sequences. Kılıc [4] gave permanent representation of Fibonacci and Lucas p-Numbers. Kılıc [6] studied on permanents and determinants of Hessenberg matrices. Yılmaz and Bozkurt [18] derived some relationships between Pell sequences, and permanents and determinants of a type of Hessenberg matrices.

In this paper, we give some determinantal and permanental representations of $F_{p,n}(x,y)$ by using various Hessenberg matrices. These results are general form of determinantal and permanental representations of polynomials and sequences mentioned Remark 1.1.

2. The determinantal representations

In this section, we give some determinantal representations of $F_{p,n}(x,y)$ by using various Hessenberg matrices.

Definition 2.1. An $n \times n$ matrix $A_n = (a_{ij})$ is called lower Hessenberg matrix if $a_{ij} = 0$ when $j - i > 1$ i.e.,

$$A_n = \begin{bmatrix} a_{11} & a_{12} & 0 & \cdots & 0 \\ a_{21} & a_{22} & a_{23} & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdots & a_{n-1,n} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdots & a_{n,n} \end{bmatrix}$$

Theorem 2.2. *[1] Let A_n be an $n \times n$ lower Hessenberg matrix for all $n \geq 1$ and $\det(A_0) = 1$. Then,*

$$\det(A_1) = a_{11}$$

and for $n \geq 2$

$$\det(A_n) = a_{n,n}\det(A_{n-1}) + \sum_{r=1}^{n-1}\left[(-1)^{n-r}a_{n,r}(\prod_{j=r}^{n-1}a_{j,j+1})\det(A_{r-1})\right].$$

Theorem 2.3. *Let $F_{p,n}(x,y)$ be the generalized bivariate Fibonacci p-polynomials and $W_{p,n} = (w_{ij})$ an $n \times n$ Hessenberg matrix defined by*

$$w_{ij} = \begin{cases} i & \text{if } i = j-1 \ , \\ x & \text{if } i = j \ , \\ i^p y & \text{if } p = i - j, \\ 0 & \text{otherwise} \end{cases}$$

that is

$$W_{p,n} = \begin{bmatrix} x & i & 0 & \cdots & 0 \\ 0 & x & i & \ddots & \vdots \\ \vdots & 0 & x & & 0 \\ i^p y & 0 & \vdots & \cdots & \\ 0 & i^p y & 0 & & 0 \\ \vdots & 0 & \ddots & x & i \\ 0 & 0 & \cdots & 0 & x \end{bmatrix}. \tag{3}$$

Then

$$\det(W_{p,n}) = F_{p,n+1}(x,y) \tag{4}$$

where $n \geq 1$ and $i = \sqrt{-1}$.

Proof. To prove (4), we use the mathematical induction on m. The result is true for $m = 1$ by hypothesis.

Assume that it is true for all positive integers less than or equal to m, namely $\det(W_{p,m}) = F_{p,m}(x, y)$. Then, we have

$$\begin{aligned}
\det(W_{p,m+1}) &= q_{m+1,m+1}\det(W_{p,m}) + \sum_{r=1}^{m}\left[(-1)^{m+1-r}q_{m+1,r}(\prod_{j=r}^{m}q_{j,j+1})\det(W_{p,r-1})\right]\\
&= x\det(W_{p,m}) + \sum_{r=1}^{m-p}\left[(-1)^{m+1-r}q_{m+1,r}(\prod_{j=r}^{m}q_{j,j+1})\det(W_{p,r-1})\right]\\
&\quad + \sum_{r=m-p+1}^{m}\left[(-1)^{m+1-r}q_{m+1,r}(\prod_{j=r}^{m}q_{j,j+1})\det(W_{p,r-1})\right]\\
&= x\det(W_{p,m}) + \left[(-1)^{p}(i)^{p}y\prod_{j=m-p+1}^{m} i\det(W_{p,m-p})\right]\\
&= x\det(W_{p,m}) + [(-1)^{p}y(i)^{p}.(i)^{p}\det(W_{p,m-p})]\\
&= x\det(W_{p,m}) + y\det(W_{p,m-p})
\end{aligned}$$

by using Theorem 2.2. From the hypothesis of induction and the definition of $F_{p,n}(x, y)$ we obtain

$$\det(W_{p,m+1}) = xF_{p,m+1}(x, y) + yF_{p,m-p+1}(x, y) = F_{p,m+2}(x, y).$$

Therefore, (4) holds for all positive integers n. □

Example 2.4. *We obtain* 6*-th* $F_{p,n}(x, y)$ *for* $p = 4$*, by using Theorem 2.3*

$$F_{4,6}(x,y) = \det\begin{bmatrix} x & i & 0 & 0 & 0\\ 0 & x & i & 0 & 0\\ 0 & 0 & x & i & 0\\ 0 & 0 & 0 & x & i\\ i^4y & 0 & 0 & 0 & x \end{bmatrix} = y + x^5.$$

Theorem 2.5. *Let* $p \geq 1$ *be an integer,* $F_{p,n}(x, y)$ *be the generalized bivariate Fibonacci p-polynomials and* $M_{p,n} = (m_{ij})$ *an* $n \times n$ *Hessenberg matrix defined by*

$$m_{ij} = \begin{cases} -1 & \textit{if } j = i+1,\\ x & \textit{if } i = j,\\ y & \textit{if } i - j = p,\\ 0 & \textit{otherwise} \end{cases}$$

that is

$$M_{p,n}=\begin{bmatrix} x & -1 & 0 & \cdots & 0 \\ 0 & x & -1 & \cdots & 0 \\ 0 & 0 & x & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ y & 0 & 0 & \cdots & 0 \\ 0 & y & 0 & \cdots & 0 \\ & \vdots & \vdots & \ddots & -1 \\ 0 & 0 & \cdots & 0 & x \end{bmatrix}. \qquad (5)$$

Then

$$\det(M_{p,n})=F_{p,n+1}(x,y).$$

Proof. Since the proof is similar to the proof of Theorem 2.3 by using Theorem 2.2 we omit the detail. □

Example 2.6. *We obtain* 6*-th* $F_{p,n}(x,y)$ *for* $p=3$*, by using Theorem 2.5*

$$F_{3,6}(x,y)=\det\begin{bmatrix} x & -1 & 0 & 0 & 0 \\ 0 & x & -1 & 0 & 0 \\ 0 & 0 & x & -1 & 0 \\ y & 0 & 0 & x & -1 \\ 0 & y & 0 & 0 & x \end{bmatrix}=2xy+x^5.$$

3. The permanent representations

Let $A=(a_{i,j})$ be a square matrix of order n over a ring R. The permanent of A is defined by

$$\mathrm{per}(A)=\sum_{\sigma\in S_n}\prod_{i=1}^{n}a_{i,\sigma(i)}$$

where S_n denotes the symmetric group on n letters.

Theorem 3.1. *[15] Let* A_n *be an* $n\times n$ *lower Hessenberg matrix for all* $n\geq 1$ *and* $per(A_0)=1$*. Then,* $per(A_1)=a_{11}$ *and for* $n\geq 2$

$$per(A_n)=a_{n,n}per(A_{n-1})+\sum_{r=1}^{n-1}\left[a_{n,r}(\prod_{j=r}^{n-1}a_{j,j+1})per(A_{r-1})\right].$$

Theorem 3.2. *Let $p \geq 1$ be an integer, $F_{p,n}(x,y)$ be the generalized bivariate Fibonacci p-polynomials and $H_{p,n} = (h_{rs})$ be an $n \times n$ lower Hessenberg matrix such that*

$$h_{rs} = \begin{cases} -i & \textit{if} \;\; s - r = 1 \;, \\ x & \textit{if} \;\; r = s \;, \\ i^p y & \textit{if} \;\; p = r - s \\ 0 & \textit{otherwise} \end{cases}$$

then

$$per(H_{p,n}) = F_{p,n+1}(x,y)$$

where $n \geq 1$ and $i = \sqrt{-1}$.

Proof. This is similar to the proof of Theorem 2.3 using Theorem 3.1. □

Example 3.3. *We obtain 6-th $F_{p,n}(x,y)$ for $p = 3$, by using Theorem 3.2*

$$F_{3,6}(x,y) = per \begin{bmatrix} x & -i & 0 & 0 & 0 \\ 0 & x & -i & 0 & 0 \\ 0 & 0 & x & -i & 0 \\ -iy & 0 & 0 & x & -i \\ 0 & -iy & 0 & 0 & x \end{bmatrix} = 2xy + x^5.$$

Theorem 3.4. *Let $p \geq 1$ be an integer, $F_{p,n}(x,y)$ be the generalized bivariate Fibonacci p-polynomials and $K_{p,n} = (k_{ij})$ be an $n \times n$ lower Hessenberg matrix such that*

$$k_{ij} = \begin{cases} 1 & \textit{if} \;\; j = i + 1, \\ x & \textit{if} \;\; i = j, \\ y & \textit{if} \;\; i - j = p, \\ 0 & \textit{otherwise.} \end{cases}$$

Then

$$per(K_{p,n}) = F_{p,n+1}(x,y).$$

Proof. This is similar to the proof of Theorem 2.3 by using Theorem 3.1. □

We note that the theorems given above are still valid for the sequences and polynomials mentioned Remark 1.1

Corollary 3.5. *If we rewrite Theorem 2.3, Theorem 2.5, Theorem 3.2 and Theorem 3.4 for x, y, p we obtain the following table.*

For	**x**	**y**	**p**	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_{p,n+1}(\mathbf{x},\mathbf{y})$,
for	x	y	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_n(\mathbf{x},\mathbf{y})$,
for	x	1	p	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_{p,n}(\mathbf{x})$,
for	x	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = f_n(x)$,
for	1	1	p	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_p(\mathbf{n})$,
for	1	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_n$,
for	$2x$	y	p	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_{p,n}(\mathbf{2x},\mathbf{y})$,
for	$2x$	y	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_n(\mathbf{2x},\mathbf{y})$,
for	$2x$	1	p	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{P}_{p,n}(\mathbf{x})$,
for	$2x$	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{P}_n(\mathbf{x})$,
for	2	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{P}_n$,
for	$2x$	-1	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{U}_{n-1}(\mathbf{x})$,
for	x	$2y$	p	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_{p,n}(\mathbf{x},\mathbf{2y})$,
for	x	$2y$	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{F}_n(\mathbf{x},\mathbf{2y})$,
for	1	$2y$	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = \mathbf{J}_n(\mathbf{y})$,
for	1	2	1	$\det(W_{p,n}) = \det(M_{p,n}) = per(H_{p,n}) = per(K_{p,n}) = J_n$.

References

[1] N.D. Cahill, J.R. D'Errico, D.A. Narayan, J.Y. Narayan, Fibonacci determinants, College Math. J. 33(2002), 221-225.

[2] M.C. Er, Sums of Fibonacci numbers by matrix method, Fibonacci Quarterly, 23(1984), 204-207.

[3] A. F. Horadam and Bro. J. M. Mahon, Pell and Pell-Lucas Polynomials, Fibonacci Quarterly, 23,1(1985), 17-20.

[4] E. Kilic and A.P. Stakhov, On the Fibonacci and Lucas p-numbers, their sums, families of bipartite graphs and permanents of certain matrices, Chaos, Solition and Fractals,(2007)

[5] E. Kilic, The Binet formula, Sums and representations of generalized Fibonacci p-numbers, Europen Journal of Combinatorics, 29(2008), 701-711.

[6] E. Kılıç and D. Tasci, On the generalized Fibonacci and Pell sequences by Hessenberg matrices, Ars Combinatoria, 94(2010) 161-174.

[7] G.-Y. Lee and S.-G. Lee, A note on generalized Fibonacci numbers, Fib. Quart., 33(1995), 273-278.

[8] G.-Y. Lee, k-Lucas numbers and associated bipartite graphs, Lineer Algebra Appl. 320(2000), 51-61.

[9] A. Lupas, A Guide of Fibonacci and Lucas Polynomial, Octagon Mathematics Magazine, 7(1999), 2-12.

[10] T. MacHenry, A Subgroup of The Group of Units in The Ring of Arithmetic Fonctions. Rocky Mount. Jour. of Math. 39(1999), 1055-1065.

[11] T. MacHenry, Generalized Fibonacci and Lucas Polynomials and Multiplicative Arithmetic Functions. Fibonacci Quarterly. 38(2000), 17-24.

[12] T. MacHenry and K. Wong, Degree k Linear Recursions mod(p) and Number Fields. Rocky Mount. Jour. of Math. *to appear.*

[13] E.P. Miles, Generalized Fibonacci Numbers and Associated Matrices. Amer. Math. Monthly 67(1960), 745-752.

[14] H. Minc, Encyclopaedia of Mathematics and its Applications, Permanents, Vol.6, Addison-Wesley Publishing Company, London, 1978.

[15] A.A. Öcal, N. Tuglu, E. Altinisik, On the representation of k-generalized Fibonacci and Lucas Numbers. Appl. Math. and Comput. 170(2005) 584-596.

[16] N. Tuglu, E.G. Kocer, A. Stakhov, Bivariate fibonacci like p-polynomials. Appl. Math. and Comput. 217(2011) 10239-10246.

[17] G. Udrea, Chebshev Polynomials and Some Methods of Approximation, Portugaliae Mathematica, Fasc.3, 55(1998), 261-269.

[18] F. Yılmaz, D. Bozkurt, Hessenberg matrices and the Pell and Perrin numbers, Journal of Number Theory. 131(2011), 1390-1396.